MycoKeys 91: 25–47 (2022)
doi: 10.3897/mycokeys.91.84970
https://mycokeys.pensoft.net

RESEARCH ARTICLE

A peer-reviewed open-access journal
MycoKeys
Launched to accelerate biodiversity research

Four new species of *Diaporthe* (Diaporthaceae, Diaporthales) from forest plants in China

Lingxue Cao[1,2,3], Dun Luo[4], Wu Lin[4], Qin Yang[1,2,3*], Xiaojun Deng[5*]

1 *Key Laboratory for Non-Wood Forest Cultivation and Conservation of the Ministry of Education, Central South University of Forestry and Technology, Changsha 410004, China* **2** *Key Laboratory of National Forestry and Grassland Administration for Control of Diseases and Pests of South Plantation, Central South University of Forestry and Technology, Changsha 410004, China* **3** *Hunan Provincial Key Laboratory for Control of Forest Diseases and Pests, Central South University of Forestry and Technology, Changsha 410004, China* **4** *Guangxi State-owned Bobai Forest Farm, Yulin, Guangxi 537600, China* **5** *Guangxi Zhuang Autonomous Region Forestry Research Institute, Nanning 530002, China*

Corresponding authors: Qin Yang (T20192466@csuft.edu.cn), Xiaojun Deng (dengxiaojun2008@sina.com)

Academic editor: Rungtiwa Phookamsak | Received 6 April 2022 | Accepted 13 June 2022 | Published 6 July 2022

Citation: Cao L, Luo D, Lin W, Yang Q, Deng X (2022) Four new species of *Diaporthe* (Diaporthaceae, Diaporthales) from forest plants in China. MycoKeys 91: 25–47. https://doi.org/10.3897/mycokeys.91.84970

Abstract

Species of *Diaporthe* inhabit a wide range of plant hosts as plant pathogens, endophytes and saprobes. During trips to collect forest pathogens in Beijing, Jiangxi, Shaanxi and Zhejiang Provinces in China, 16 isolates of *Diaporthe* were obtained from branch cankers and leaf spots. These isolates were studied by applying a polyphasic approach including morphological, cultural data, and phylogenetic analyses of the nuclear ribosomal internal transcribed spacer (ITS), calmodulin (*cal*), histone H3 (*his3*), partial translation elongation factor-1α (*tef-1a*) and β-tubulin (*tub2*) loci. Results revealed four new taxa, *D. celticola*, *D. meliae*, *D. quercicola*, *D. rhodomyrti* **spp. nov.** and two known species, *D. eres* and *D. multiguttulata*.

Keywords

Diaporthaceae, DNA phylogeny, four new taxa, systematics, taxonomy

* These authors contributed equally to this work.

Copyright *Lingxue Cao et al.* This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Introduction

Diaporthe Nitschkes (syn. *Phomopsis*) is a large genus in the Diaporthaceae with plant pathogens, endophytes or saprobes (Muralli et al. 2006; Rossman et al. 2007; Santos and Phillips 2009; Santos et al. 2011; Udayanga et al. 2011, 2014a, b, 2015; Fan et al. 2015, 2018; Du et al. 2016; Dissanayake et al. 2017; Guarnaccia and Crous 2017, 2018; Guarnaccia et al. 2018; Yang et al. 2018, 2020, 2021a, b; Guo et al. 2020; Sun et al. 2021). Currently, more than 1100 epithets for *Diaporthe* and 950 for *Phomopsis* are listed in Index Fungorum (http://www.indexfungorum.org/; accessed 1 April 2022) with names often based on host association.

The family Diaporthaceae was established by von Höhnel (1917) and was accommodated in the order Diaporthales. Wehmeyer (1975) confined this family to include *Diaporthe* and *Mazzantia*. Later, Diaporthaceae was synonymised under Valsaceae (Barr 1978). However, analysis of LSU sequence data of diaporthalean taxa showed the distinct placement of Diaporthaceae in Diaporthales where it formed a well-supported clade (Castlebury et al. 2002). *Diaporthe*, the type genus of Diaporthaceae, is characterised by immersed ascomata and an erumpent pseudostroma with elongated perithecial necks (Gomes et al. 2013). Asci are unitunicate, clavate to cylindrical. Ascospores are fusoid, ellipsoid to cylindrical, hyaline, biseriate to uniseriate in the ascus, sometimes with appendages (Udayanga et al. 2011). The asexual morph is characterised by ostiolate conidiomata, with cylindrical phialides producing three types (alpha, beta, and gamma conidia) of hyaline, aseptate conidia (Udayanga et al. 2011; Gomes et al. 2013).

In China, the classification of *Diaporthe* has been progressing and the basis for the species identification is a combination of morphological, cultural and phylogenetical analyses (Huang et al. 2015; Gao et al. 2017; Guarnaccia and Crous 2017; Yang et al. 2017, 2018, 2020, 2021a, b; Manawasinghe et al. 2019; Jiang et al. 2021; Huang et al. 2021; Sun et al. 2021; Wang et al. 2021). The present study was conducted to identify *Diaporthe* species that cause dieback and leaf spot disease in Beijing, Jiangxi, Shaanxi and Zhejiang Provinces based on modern taxonomic concepts.

Materials and methods

Fungal isolation

From 2018 to 2020, sample collections have been ongoing in Beijing, Jiangxi, Shaanxi and Zhejiang Provinces, China (Table 1). Collected samples were taken to the laboratory for isolation and photographed, documented and then kept at 4 °C for further study.

A total of 16 isolates from host material were obtained by removing a mucoid conidia mass from conidiomata, spreading the suspension on the surface of 1.8% potato dextrose agar (PDA), and incubating at 25 °C for up to 24 h. Single germinating conidium was removed and plated onto fresh PDA plates. Specimens were deposited in the Museums of the Beijing Forestry University (BJFC) and Central South University

Table 1. Isolates and GenBank accession numbers of sequences used in this study.

Species	Isolate	GenBank accession numbers				
		ITS	*cal*	*his3*	*tef-1α*	*tub2*
Diaporthe acaciigena	CBS 129521*	KC343005	KC343247	KC343489	KC343731	KC343973
Diaporthe acericola	MFLUCC 17-0956*	KY964224	KY964137	NA	KY964180	KY964074
Diaporthe acerigena	CFCC 52554*	MH121489	MH121413	MH121449	MH121531	NA
Diaporthe acerigena	CFCC 52555	MH121490	MH121414	MH121450	MH121532	NA
Diaporthe acuta	PSCG 047*	MK626957	MK691125	MK726161	MK654802	MK691225
Diaporthe acutispora	LC6161*	KX986764	KX999274	KX999235	KX999155	KX999195
Diaporthe alangii	CFCC 52556*	MH121491	MH121415	MH121451	MH121533	MH121573
Diaporthe alangii	CFCC 52557	MH121492	MH121416	MH121452	MH121534	MH121574
Diaporthe albosinensis	CFCC 53066*	MK432659	MK442979	MK443004	MK578133	MK578059
Diaporthe albosinensis	CFCC 53067	MK432660	MK442980	MK443005	MK578134	MK578060
Diaporthe alleghaniensis	CBS 495.72*	KC343007	KC343249	KC343491	KC343733	KC343975
Diaporthe ambigua	CBS 114015*	KC343010	KC343252	KC343494	KC343736	KC343978
Diaporthe ampelina	STE-U 2660	AF230751	AY745026	NA	AY745056	JX275452
Diaporthe amygdali	CBS 126679*	MH864208	KC343264	KC343506	KC343748	KC343990
Diaporthe amygdali syn. *D. chongqingensis*	PSCG 435	MK626916	MK691209	MK726257	MK654866	MK691321
Diaporthe amygdali syn. *D. fusicola*	CGMCC 3.17087	KF576281	KF576233	NA	KF576256	KF576305
Diaporthe amygdali syn. *D. garethjonesii*	MFLUCC 12-0542a	KT459423	KT459470	NA	KT459457	KT459441
Diaporthe amygdali syn. *D. kadsurae*	CFCC 52586	MH121521	MH121439	MH121479	MH121563	MH121600
Diaporthe amygdali syn. *D. kadsurae*	CFCC 52587	MH121522	MH121440	MH121480	MH121564	MH121601
Diaporthe amygdali syn. *D. mediterranea*	SAUCC194.111	MT822639	MT855718	MT855606	MT855836	MT855951
Diaporthe amygdali syn. *D. ovoicicola*	CGMCC 3.17093	KF576265	KF576223	NA	KF576240	KF576289
Diaporthe amygdali syn. *D. sterilis*	CBS 136969	KJ160579	KJ160548	MF418350	KJ160611	KJ160528
Diaporthe amygdali syn. *D. ternstroemiae*	CGMCC 3.15183	KC153098	NA	NA	KC153089	NA
Diaporthe anacardii	CBS 720.97*	KC343024	KC343266	KC343508	KC343750	KC343992
Diaporthe angelicae	CBS 111592*	KC343027	KC343269	KC343511	KC343753	KC343995
Diaporthe apiculata	CFCC 53068	MK432651	MK442973	MK442998	MK578127	MK578054
Diaporthe apiculata	CFCC 53069	MK432652	MK442974	MK442999	MK578128	MK578055
Diaporthe aquatic	IFRDCC 3051*	JQ797437	NA	NA	NA	NA
Diaporthe arctii	DP0482*	KJ590736	KJ612133	KJ659218	KJ590776	KJ610891
Diaporthe arecae	CBS 161.64*	KC343032	KC343274	KC343516	KC343758	KC344000
Diaporthe arengae	CBS 114979*	KC343034	KC343276	KC343518	KC343760	KC344002
Diaporthe arezzoensis	MFLUCC 15-0127*	MT185503	NA	NA	NA	NA
Diaporthe aseana	MFLUCC 12-0299a*	KT459414	KT459464	NA	KT459448	KT459432
Diaporthe asheicola	CBS 136967*	KJ160562	KJ160542	NA	KJ160594	KJ160518
Diaporthe aspalathi	CBS 117169*	KC343036	KC343278	KC343520	KC343762	KC344004
Diaporthe australafricana	CBS 111886*	KC343038	KC343280	KC343522	KC343764	KC344006
Diaporthe australiana	CBS 146457*	MN708222	NA	NA	MN696522	MN696530
Diaporthe baccae	CBS 136972*	KJ160565	MG281695	MF418264	KJ160597	MF418509
Diaporthe batatas	CBS 122.21	KC343040	KC343282	KC343524	KC343766	KC344008
Diaporthe bauhiniae	CFCC 53071*	MK432648	MK442970	MK442995	MK578124	MK578051
Diaporthe bauhiniae	CFCC 53072	MK432649	MK442971	MK442996	MK578125	MK578052
Diaporthe beilharziae	BRIP 54792*	JX862529	NA	NA	JX862535	KF170921
Diaporthe benedicti	SBen914*	KM669929	KM669862	NA	KM669785	NA
Diaporthe betulae	CFCC 50469*	KT732950	KT732997	KT732999	KT733016	KT733020
Diaporthe betulae	CFCC 50470	KT732951	KT732998	KT733000	KT733017	KT733021
Diaporthe betulicola	CFCC 51128*	KX024653	KX024659	KX024661	KX024655	KX024657
Diaporthe betulicola	CFCC 51129	KX024654	KX024660	KX024662	KX024656	KX024658
Diaporthe betulina	CFCC 52560*	MH121495	MH121419	MH121455	MH121537	MH121577
Diaporthe betulina	CFCC 52561	MH121496	MH121420	MH121456	MH121538	MH121578
Diaporthe biconispora	ZJUD62*	KJ490597	NA	KJ490539	KJ490476	KJ490418
Diaporthe biguttulata	ZJUD47*	KJ490582	NA	KJ490524	KJ490461	KJ490403
Diaporthe bohemiae	CBS 143347*	MG281015	MG281710	MG281361	MG281536	MG281188
Diaporthe brasiliensis	CBS 133183*	KC343042	KC343284	KC343526	KC343768	KC344010
Diaporthe caatingaensis	URM7486*	KY085927	KY115597	KY115605	KY115603	KY115600
Diaporthe camelliae-sinensis	SAUCC194.92*	MT822620	MT855699	MT855588	MT855932	MT855817
Diaporthe canthi	CPC 19740*	JX069864	KC843174	NA	KC843120	KC843230
Diaporthe caryae	CFCC 52563*	MH121498	MH121422	MH121458	MH121540	MH121580
Diaporthe caryae	CFCC 52564	MH121499	MH121423	MH121459	MH121541	MH121581
Diaporthe cassines	CPC 21916*	KF777155	NA	NA	KF777244	NA

Species	Isolate	GenBank accession numbers				
		ITS	*cal*	*his3*	*tef-1α*	*tub2*
Diaporthe caulivora	CBS 127268*	MH864501	KC343287	KC343529	KC343771	KC344013
Diaporthe celticola	**CFCC 53074***	**MK573948**	**MK574587**	**MK574603**	**MK574623**	**MK574643**
Diaporthe celticola	**CFCC 53075**	**MK573949**	**MK574588**	**MK574604**	**MK574624**	**MK574644**
Diaporthe celticola	**CFCC 53076**	**MK573950**	**MK574589**	**MK574605**	**MK574625**	**MK574645**
Diaporthe cercidis	CFCC 52565*	MH121500	MH121424	MH121460	MH121542	MH121582
Diaporthe cercidis	CFCC 52566	MH121501	MH121425	MH121461	MH121543	MH121583
Diaporthe chamaeropis	CBS 454.81*	KC343048	KC343290	KC343532	KC343774	KC344016
Diaporthe charlesworthii	BRIP 54884m*	KJ197288	NA	NA	KJ197250	KJ197268
Diaporthe chensiensis	CFCC 52567*	MH121502	MH121426	MH121462	MH121544	MH121584
Diaporthe chensiensis	CFCC 52568	MH121503	MH121427	MH121463	MH121545	MH121585
Diaporthe chrysalidocarpi	SAUCC194.35*	MT822563	MT855646	MT855532	MT855760	MT855876
Diaporthe cichorii	MFLUCC 17-1023*	KY964220	KY964133	NA	KY964176	KY964104
Diaporthe cinnamomi	CFCC 52569*	MH121504	NA	MH121464	MH121546	MH121586
Diaporthe cinnamomi	CFCC 52570	MH121505	NA	MH121465	MH121547	MH121587
Diaporthe cissampeli	CPC 27302*	KX228273	NA	KX228366	NA	KX228384
Diaporthe citri	AR3405*	KC843311	KC843157	KJ420881	KC843071	KC843187
Diaporthe citri	CFCC 53079	MK573940	MK574579	MK574595	MK574615	MK574635
Diaporthe citriasiana	CGMCC 3.15224*	JQ954645	KC357491	KJ490515	JQ954663	KC357459
Diaporthe citrichinensis	CGMCC 3.15225*	JQ954648	KC357494	KJ420880	JQ954666	KJ490396
Diaporthe collariana	MFLU 17-2770*	MG806115	MG783042	NA	MG783040	MG783041
Diaporthe compactum	LC3083*	KP267854	NA	KP293508	KP267928	KP293434
Diaporthe conica	CFCC 52571*	MH121506	MH121428	MH121466	MH121548	MH121588
Diaporthe conica	CFCC 52572	MH121507	MH121429	MH121467	MH121549	MH121589
Diaporthe constrictospora	CGMCC 3.20096*	MT385947	MT424718	MW022487	MT424682	MT424702
Diaporthe convolvuli	CBS 124654	KC343054	KC343296	KC343538	KC343780	KC344022
Diaporthe coryli	CFCC 53083*	MK432661	MK442981	MK443006	MK578135	MK578061
Diaporthe coryli	CFCC 53084	MK432662	MK442982	MK443007	MK538176	MK578062
Diaporthe corylicola	CFCC 53986*	MW839880	MW836684	MW836717	MW815894	MW883977
Diaporthe corylicola	CFCC 53987	MW839867	MW836685	MW836718	MW815895	MW883978
Diaporthe crotalariae	CBS 162.33*	MH855395	JX197439	KC343540	GQ250307	KC344024
Diaporthe crousii	CAA 823*	MK792311	MK883835	MK871450	MK828081	MK837932
Diaporthe cucurbitae	DAOM 42078*	KM453210	NA	KM453212	KM453211	KP118848
Diaporthe cuppatea	CBS 117499	MH863021	KC343299	KC343541	KC343783	KC344025
Diaporthe cynaroidis	CBS 122676*	KC343058	KC343300	KC343542	KC343784	KC344026
Diaporthe cytosporella	FAU461	KC843307	KC843141	MF418283	KC843116	KC843221
Diaporthe diospyricola	CPC 21169*	KF777156	NA	NA	NA	NA
Diaporthe discoidispora	ZJUD89*	KJ490624	NA	KJ490566	KJ490503	KJ490445
Diaporthe dorycnii	MFLUCC 17-1015*	KY964215	NA	NA	KY964171	KY964099
Diaporthe drenthii	CBS 146453*	MN708229	NA	NA	MN696526	MN696537
Diaporthe durionigena	VTCC 930005*	MN453530	NA	NA	MT276157	MT276159
Diaporthe elaeagni-glabrae	LC4802*	KX986779	KX999281	KX999251	KX999171	KX999212
Diaporthe endophytica	CBS 133811*	KC343065	KC343307	KC343549	KC343791	KC344033
Diaporthe eres	AR5193*	KJ210529	KJ434999	KJ420850	KJ210550	KJ420799
Diaporthe eres	AR5211	KJ210538	KJ435043	KJ420875	KJ210559	KJ420828
Diaporthe eres	CBS 587.79	KC343153	KC343395	KC343637	KC343879	KC344121
Diaporthe eres	CFCC 52575	MH121510	NA	MH121470	MH121552	MH121592
Diaporthe eres	CFCC 52576	MH121511	MH121432	MH121471	MH121553	MH121593
Diaporthe eres	CFCC 52577	MH121512	MH121433	MH121472	MH121554	MH121594
Diaporthe eres	CFCC 52578	MH121513	MH121434	MH121473	MH121555	MH121595
Diaporthe eres	CFCC 52579	MH121514	NA	MH121474	MH121556	NA
Diaporthe eres	CFCC 52580	MH121515	NA	MH121475	MH121557	MH121596
Diaporthe eres	CFCC 52581	MH121516	NA	MH121476	MH121558	MH121597
Diaporthe eres	CGMCC 3.15181	KC153096	NA	NA	KC153087	KF576312
Diaporthe eres	CGMCC 3.17081	KF576282	NA	NA	KF576257	KF576306
Diaporthe eres	CGMCC 3.17089	KF576267	NA	NA	KF576242	KF576291
Diaporthe eres	DAOM 695742	KU552025	NA	NA	KU552023	KU574615
Diaporthe eres	MAFF 625034	JQ807469	KJ435023	KJ420868	JQ807418	KJ420819
Diaporthe eres	MFLU 17-0646	MG828895	MG829274	NA	MG829270	MG843877
Diaporthe eres	MFLUCC 16-0113	KU557563	KU557611	NA	KU557631	KU557587
Diaporthe eres	MFLUCC 17-0963	KY964190	KY964116	NA	KY964146	KY964073
Diaporthe eres syn. *D. alnea*	CBS 146.46	KC343008	KC343250	KC343492	KC343734	KC343976

Species	Isolate	GenBank accession numbers				
		ITS	*cal*	*his3*	*tef-1α*	*tub2*
Diaporthe eres syn. *D. camptothecicola*	CFCC 51632	KY203726	KY228877	KY228881	KY228887	KY228893
Diaporthe eres syn. *D. celastrina*	CBS 139.27	KC343047	KC343289	KC343531	KC343773	KC344015
Diaporthe eres syn. *D. celeris*	CBS 143349	MG281017	MG281712	MG281363	MG281538	MG281190
Diaporthe eres syn.*D. ellipicola*	CGMCC 3.17084	KF576270	NA	NA	KF576245	KF576294
Diaporthe eres syn. D. neilliae	CBS 144.27	KC343144	KC343386	KC343628	KC343870	KC344112
Diaporthe eres syn. *D. pulla*	CBS 338.89	KC343152	KC343394	KC343636	KC343878	KC344120
Diaporthe eres	**CSUFTCC101**	**ON076564**	**NA**	**ON081664**	**ON081656**	**NA**
Diaporthe eres	**CSUFTCC102**	**ON076565**	**NA**	**ON081665**	**ON081657**	**NA**
Diaporthe eres	**CSUFTCC103**	**ON076566**	**NA**	**ON081666**	**ON081658**	**NA**
Diaporthe eucalyptorum	CBS 132525*	MH305525	NA	NA	NA	NA
Diaporthe foeniculacea	CBS 111553*	KC343101	KC343343	KC343585	KC343827	KC344069
Diaporthe fraxini-angustifoliae	BRIP 54781*	JX862528	NA	NA	JX862534	KF170920
Diaporthe fraxinicola	CFCC 52582*	MH121517	MH121435	NA	MH121559	NA
Diaporthe fraxinicola	CFCC 52583	MH121518	MH121436	NA	MH121560	NA
Diaporthe fructicola	MAFF 246408*	LC342734	LC342738	LC342737	LC342735	LC342736
Diaporthe fulvicolor	PSCG 051*	MK626859	MK691132	MK726163	MK654806	MK691236
Diaporthe ganjae	CBS 180.91*	KC343112	KC343354	KC343596	KC343838	KC344080
Diaporthe ganzhouensis	CFCC 53087*	MK432665	MK442985	MK443010	MK578139	MK578065
Diaporthe ganzhouensis	CFCC 53088	MK432666	MK442986	MK443011	MK578140	MK578066
Diaporthe goulteri	BRIP 55657a*	KJ197290	NA	NA	KJ197252	KJ197270
Diaporthe grandiflori	SAUCC194.84*	MT822612	MT855691	MT855580	MT855809	MT855924
Diaporthe guangxiensis	JZB320087	MK335765	MK736720	NA	MK500161	MK523560
Diaporthe gulyae	BRIP 54025	JF431299	NA	NA	JN645803	KJ197271
Diaporthe guttulata	CGMCC 3.20100*	MT385950	MW022470	MW022491	MT424685	MT424705
Diaporthe helianthi	CBS 592.81*	KC343115	KC343357	KC343599	KC343841	KC344083
Diaporthe heliconiae	SAUCC194.77*	MT822605	MT855684	MT855573	MT855802	MT855917
Diaporthe heterophyllae	CPC 26215*	MG600222	MG600218	MG600220	MG600224	MG600226
Diaporthe heterostemmatis	SAUCC194.85*	MT822613	MT855692	MT855581	MT855810	MT855925
Diaporthe hickoriae	CBS 145.26*	KC343118	KC343360	KC343620	KC343844	KC344086
Diaporthe hispaniae	CBS 143351*	MG281123	MG281820	MG281471	MG281644	MG281296
Diaporthe hongkongensis	CBS 115448*	KC343119	KC343361	KC343603	KC343845	KC344087
Diaporthe hubeiensis	JZB320123*	MK335809	MK500235	NA	MK523570	MK500148
Diaporthe incomplete	LC6754*	KX986794	KX999289	KX999265	KX999186	KX999226
Diaporthe inconspicua	CBS 133813*	KC343123	KC343365	KC343607	KC343849	KC344091
Diaporthe infecunda	CBS 133812*	KC343126	KC343368	KC343610	KC343852	KC344094
Diaporthe irregularis	CGMCC 3.20092*	MT385951	MT424721	NA	MT424686	MT424706
Diaporthe isoberliniae	CPC 22549*	KJ869190	NA	NA	NA	KJ869245
Diaporthe juglandicola	CFCC 51134*	KU985101	KX024616	KX024622	KX024628	KX024634
Diaporthe kochmanii	BRIP 54033*	JF431295	NA	NA	JN645809	NA
Diaporthe kongii	BRIP 54031*	JF431301	NA	NA	JN645797	KJ197272
Diaporthe krabiensis	MFLUCC 17-2481*	MN047100	NA	NA	MN433215	MN431495
Diaporthe lenispora	CGMCC 3.20101*	MT385952	MW022472	MW022493	MT424687	MT424707
Diaporthe litchicola	BRIP 54900*	JX862533	NA	NA	JX862539	KF170925
Diaporthe litchi	SAUCC194.22*	MT822550	MT855635	MT855519	MT855747	MT855863
Diaporthe lithocarpi	CGMCC 3.15175*	KC135104	KF576235	NA	KC153095	KF576311
Diaporthe longicolla	FAU599	KJ590728	KJ612124	KJ659188	KJ590767	KJ610883
Diaporthe longispora	CBS 194.36*	MH855769	KC343377	KC343619	KC343861	KC344103
Diaporthe lusitanicae	CBS 123212*	MH863279	KC343378	KC343620	KC343862	KC344104
Diaporthe lutescens	SAUCC194.36*	MT822564	MT855647	MT855533	MT855761	MT855877
Diaporthe macadamiae	CBS 146455*	MN708230	NA	NA	MN696528	MN696539
Diaporthe macintoshii	BRIP 55064a*	KJ197289	NA	NA	KJ197251	KJ197269
Diaporthe malorum	CAA 734*	KY435638	KY435658	KY435648	KY435627	KY435668
Diaporthe marina	MFLU 17-2622*	MN047102	NA	NA	NA	NA
Diaporthe masirevicii	BRIP 54256*	KJ197276	NA	NA	KJ197238	KJ197256
Diaporthe mayteni	CBS 133185*	KC343139	KC343381	KC343623	KC343865	KC344107
Diaporthe maytenicola	CPC 21896*	KF777157	NA	NA	NA	KF777250
Diaporthe melastomatis	SAUCC194.55*	MT822583	MT855664	MT855551	MT855780	MT855896
Diaporthe melonis	CBS 435.87	KC343141	KC343383	KC343625	KC343867	KC344109
Diaporthe meliae	**CFCC 53089***	**MK432657**	**NA**	**ON081662**	**ON081654**	**MK578057**
Diaporthe meliae	**CFCC 53090**	**MK432658**	**NA**	**ON081663**	**ON081655**	**MK578058**
Diaporthe middletonii	BRIP 54884e*	KJ197286	NA	NA	KJ197248	KJ197266

Species	Isolate	GenBank accession numbers				
		ITS	*cal*	*his3*	*tef-1α*	*tub2*
Diaporthe minima	CGMCC 3.20097*	MT385953	MT424722	MW022496	MT424688	MT424708
Diaporthe minusculata	CGMCC 3.20098*	MT385957	MW022475	MW022499	MT424692	MT424712
Diaporthe miriciae	BRIP 54736j*	KJ197282	NA	NA	KJ197244	KJ197262
Diaporthe multigutullata	CFCC 53095	MK432645	MK442967	MK442992	MK578121	MK578048
Diaporthe multigutullata	CFCC 53096	MK432646	MK442968	MK442993	MK578122	MK578049
Diaporthe multigutullata	**CFCC 53098**	**MK573957**	**MK574592**	**MK574612**	**MK574632**	**MK574652**
Diaporthe multigutullata	**CFCC 53099**	**MK573958**	**MK574593**	**MK574613**	**MK574633**	**MK574653**
Diaporthe multigutullata	**CFCC 53100**	**MK573959**	**MK574594**	**MK574614**	**MK574634**	**MK574654**
Diaporthe musigena	CBS 129519*	KC343143	KC343385	KC343267	KC343869	KC344111
Diaporthe myracrodruonis	URM 7972	MK205289	MK205290	NA	MK213408	MK205291
Diaporthe neoarctii	CBS 109490*	KC343145	KC343387	KC343629	KC343871	KC344113
Diaporthe neoraonikayaporum	MFLUCC 14-1136*	KU712449	KU749356	NA	KU749369	KU743988
Diaporthe nothofagi	BRIP 54801*	JX862530	NA	NA	JX862536	KF170922
Diaporthe novem	CBS 127269	KC343155	KC343397	KC343639	KC343881	KC344123
Diaporthe ocoteae	CPC 26217*	KX228293	NA	NA	NA	KX228388
Diaporthe oraccinii	LC3166*	KP267863	NA	KP293517	KP267937	KP293443
Diaporthe ovalispora	ZJUD93*	KJ490628	NA	KJ490570	KJ490507	KJ490449
Diaporthe oxe	CBS 133186*	KC343164	KC343406	KC343648	KC343890	KC344132
Diaporthe padina	CFCC 52590*	MH121525	MH121443	MH121483	MH121567	MH121604
Diaporthe padina	CFCC 52591	MH121526	MH121444	MH121484	MH121568	MH121605
Diaporthe pandanicola	MFLUCC 17-0607*	MG646974	NA	NA	NA	MG646930
Diaporthe paranensis	CBS 133184*	KC343171	KC343413	KC343655	KC343897	KC344139
Diaporthe parapterocarpi	CPC 22729	KJ869138	NA	NA	NA	KJ869248
Diaporthe parvae	PSCG 035	MK626920	MK691169	MK726211	MK654859	MK691249
Diaporthe pascoei	BRIP 54847*	JX862538	NA	NA	JX862538	KF170924
Diaporthe passiflorae	CPC 19183*	JX069860	KY435644	KY435654	KY435623	KY435674
Diaporthe passifloricola	CPC 27480*	KX228292	NA	KX228367	NA	KX228387
Diaporthe penetriteum	LC3215	KP267879	NA	KP293532	KP267953	NA
Diaporthe perjuncta	CBS 109745*	KC343172	KC343414	KC343656	KC343898	KC344140
Diaporthe perseae	CBS 151.73	KC343173	KC343415	KC343657	KC343899	KC343141
Diaporthe pescicola	MFLUCC 16-0105*	KU557555	KU557603	NA	KU557623	KU557579
Diaporthe phaseolorum	AR4203*	KJ590738	KJ612135	KJ659220	KJ590739	KJ610893
Diaporthe phillipsii	CAA 817*	MK792305	MK883831	MK871445	MK828076	MN000351
Diaporthe podocarpi-macrophylli	LC6155*	KX986774	KX999278	KX999246	KX999167	KX999207
Diaporthe pometiae	SAUCC194.72*	MT822600	MT855679	MT855568	MT855797	MT855912
Diaporthe pseudoalnea	CFCC 54190*	MZ727037	MZ753468	MZ781302	MZ816343	MZ753487
Diaporthe pseudomangiferae	CBS 101339*	KC343181	KC343423	KC343665	KC343907	KC344149
Diaporthe pseudophoenicicola	CBS 176.77	KC343183	KC343425	KC343667	KC343909	KC344151
Diaporthe pseudotsugae	MFLU 15-3228*	KY964225	KY964138	NA	KY964181	KY964108
Diaporthe psoraleae	CPC 21634*	KF777158	NA	NA	KF777245	KF777251
Diaporthe psoraleae-pinnatae	CPC 21638*	KF777159	NA	NA	NA	KF777252
Diaporthe pterocarpi	MFLUCC 10-0575*	JQ619901	JX197453	NA	JX275418	NA
Diaporthe pterocarpicola	MFLUCC 10-0580a*	JQ619887	JX197433	NA	JX275403	JX275441
Diaporthe pungensis	SAUCC194.112*	MT822640	MT855719	MT855607	MT855837	MT855952
Diaporthe pyracanthae	CAA483	KY435635	KY435656	KY435645	KY435625	KY435666
Diaporthe quercicola	**CSUFTCC104***	**ON076567**	**ON081670**	**ON081667**	**ON081659**	**NA**
Diaporthe quercicola	**CSUFTCC105**	**ON076568**	**ON081671**	**ON081668**	**ON081660**	**NA**
Diaporthe quercicola	**CSUFTCC106**	**ON076569**	**ON081672**	**ON081669**	**ON081661**	**NA**
Diaporthe racemosae	CPC 26646*	MG600223	MG600219	MG600221	MG600225	MG600227
Diaporthe raonikayaporum	CBS 133182*	KC343188	KC343430	KC343672	KC343914	KC344156
Diaporthe ravennica	MFLUCC 16-0997	NA	NA	NA	MT394670	NA
Diaporthe rhodomyrti	**CFCC 53101***	**MK432643**	**MK442965**	**MK442990**	**MK578119**	**MK578046**
Diaporthe rhodomyrti	**CFCC 53102**	**MK432644**	**MK442966**	**MK442991**	**MK578120**	**MK578047**
Diaporthe rhusicola	CPC 18191*	JF951146	KC843124	NA	KC843100	KC843205
Diaporthe rosae	MFLUCC 17-2658*	MG828894	MG829273	NA	NA	MG843878
Diaporthe rosiphthora	COAD 2914	MT311197	MT313691	NA	MT313693	NA
Diaporthe rossmaniae	CAA 762*	MK792290	MK883822	MK871432	MK828063	MK837914
Diaporthe rostrata	CFCC 50062*	KP208847	KP208849	KP208851	KP208853	KP208855
Diaporthe rostrata	CFCC 50063	KP208848	KP208850	KP208852	KP208854	KP208856
Diaporthe rudis	AR3422	KC843331	KC843146	NA	KC843090	KC843177
Diaporthe saccarata	CBS 116311*	KC343190	KC343432	KC343674	KC343916	KC344158

Species	Isolate	GenBank accession numbers				
		ITS	*cal*	*his3*	*tef-1α*	*tub2*
Diaporthe sackstonii	BRIP 54669b*	KJ197287	NA	NA	KJ197249	KJ197267
Diaporthe salicicola	BRIP 54825*	JX862531	NA	NA	JX862537	KF170923
Diaporthe sambucusii	CFCC 51986*	KY852495	KY852499	KY852503	KY852507	KY852511
Diaporthe sambucusii	CFCC 51987	KY852496	KY852500	KY852504	KY852508	KY852512
Diaporthe schimae	CFCC 53103*	MK442640	MK442962	MK442987	MK578116	MK578043
Diaporthe schimae	CFCC 53104	MK442641	MK442963	MK442988	MK578117	MK578044
Diaporthe schini	CBS 133181*	KC343191	KC343433	KC343675	KC343917	KC344159
Diaporthe schisandrae	CFCC 51988*	KY852497	KY852501	KY852505	KY852509	KY852513
Diaporthe schisandrae	CFCC 51989	KY852498	KY852502	KY852506	KY852510	KY852514
Diaporthe schoeni	MFLU 15-1279*	KY964226	KY964139	NA	KY964182	KY964109
Diaporthe sclerotioides	CBS 296.67*	MH858974	KC343435	KC343677	KC343919	KC344161
Diaporthe searlei	CBS 146456*	MN708231	NA	NA	NA	MN696540
Diaporthe sennae	CFCC 51636*	KY203724	KY228875	NA	KY228885	KY228891
Diaporthe sennae	CFCC 51637	KY203725	KY228876	NA	KY228886	KY228892
Diaporthe sennicola	CFCC 51634*	KY203722	KY228873	KY228879	KY228883	KY228889
Diaporthe sennicola	CFCC 51635	KY203723	KY228874	KY228880	KY228884	KY228890
Diaporthe serafiniae	BRIP 55665a*	KJ197274	NA	NA	KJ197236	KJ197254
Diaporthe shaanxiensis	CFCC 53106*	MK432654	MK442976	MK443001	MK578130	NA
Diaporthe shaanxiensis	CFCC 53107	MK432655	MK432977	MK432002	MK578131	NA
Diaporthe siamensis	MFLUCC 10-0573a*	JQ619879	JX197423	NA	JX275393	JX275429
Diaporthe silvicola	CFCC 54191*	MZ727041	MZ753472	MZ753481	MZ816347	MZ753491
Diaporthe sojae	FAU635	KJ590719	KJ612116	KJ659208	KJ590762	KJ610875
Diaporthe spartinicola	CPC 24951*	KR611879	NA	KR857696	NA	KR857695
Diaporthe spinosa	PSCG 383*	MK626849	MK691129	MK726156	MK654811	MK691234
Diaporthe stictica	CBS 370.54	KC343212	KC343454	KC343696	KC343938	KC344180
Diaporthe subclavata	ZJUD95*	KJ490630	NA	KJ490572	KJ490509	KJ490451
Diaporthe subcylindrospora	KUMCC 17-0151	MG746629	NA	NA	MG746630	MG746631
Diaporthe subellipicola	KUMCC 17-0153*	MG746632	NA	NA	MG746633	MG746634
Diaporthe subordinaria	CBS 464.90	KC343214	KC343456	KC343698	KC343940	KC344182
Diaporthe taoicola	MFLUCC 16-0117*	KU557567	NA	NA	KU557635	KU557591
Diaporthe tectonae	MFLUCC 12-0777*	KU712430	KU749345	NA	KU749359	KU743977
Diaporthe tectonendophytica	MFLUCC 13-0471*	KU712439	KU749354	NA	KU749367	KU743986
Diaporthe tectonigena	LC6512	KX986782	KX999284	KX999254	KX999174	KX999214
Diaporthe terebinthifolii	CBS 133180*	KC343216	KC343458	KC343700	KC343942	KC344184
Diaporthe thunbergii	MFLUCC 10-0576a*	JQ619893	JX197440	NA	JX275409	JX275449
Diaporthe thunbergiicola	MFLUCC 12-0033*	KP715097	NA	NA	KP715098	NA
Diaporthe tibetensis	CFCC 51999*	MF279843	MF279888	MF279828	MF279858	MF279873
Diaporthe tibetensis	CFCC 52000	MF279844	MF279889	MF279829	MF279859	MF279874
Diaporthe torilicola	MFLUCC 17-1051*	KY964212	KY964127	NA	KY964168	KY964096
Diaporthe toxica	CBS 534.93*	KC343220	KC343462	KC343704	KC343946	KC344188
Diaporthe tulliensis	BRIP 62248a*	KR936130	NA	NA	KR936133	KR936132
Diaporthe ueckerae	FAU656*	KJ590726	KJ612122	KJ659215	KJ590747	KJ610881
Diaporthe ukurunduensis	CFCC 52592*	MH121527	MH121445	MH121485	MH121569	NA
Diaporthe ukurunduensis	CFCC 52593	MH121528	MH121446	MH121486	MH121570	NA
Diaporthe undulate	LC6624*	KX986798	NA	KX999269	KX999190	KX999230
Diaporthe unshiuensis	CFCC 52594	MH121529	MH121447	MH121487	MH121571	MH121606
Diaporthe unshiuensis	CFCC 52595	MH121530	MH121448	MH121488	MH121572	MH121607
Diaporthe vaccinii	CBS 160.32*	KC343228	KC343470	KC343712	KC343954	KC343196
Diaporthe vangueriae	CBS 137985*	KJ869137	NA	NA	NA	KJ869247
Diaporthe vawdreyi	BRIP 57887a*	KR936126	NA	NA	KR936129	KR936128
Diaporthe velutina	LC4421	KX986790	NA	KX999261	KX999182	KX999223
Diaporthe verniciicola	CFCC 53109*	MK573944	MK574583	MK574599	MK574619	MK574639
Diaporthe verniciicola	CFCC 53110	MK573945	MK574584	MK574600	MK574620	MK574640
Diaporthe viniferae	JZB320071	MK341551	MK500119	NA	MK500107	MK500112
Diaporthe virgiliae	CMW 40748	KP247556	NA	NA	NA	KP247575
Diaporthe xishuangbanica	LC6707*	KX986783	NA	KX999255	KX999175	KX999216
Diaporthe xunwuensis	CFCC 53085*	MK432663	MK442983	MK443008	MK578137	MK578063
Diaporthe xunwuensis	CFCC 53086	MK432664	MK442984	MK443009	MK578138	MK578064
Diaporthe yunnanensis	LC6168*	KX986796	KX999290	KX999267	KX999188	KX999228
Diaporthe zaobaisu	PSCG 031*	MK626922	NA	MK726207	MK654855	MK691245
Diaporthella corylina	CBS 121124*	KC343004	KC343246	KC343488	KC343730	KC343972

Strains in this study are marked in bold. NA: Not available. Ex-type/ex-epitype isolates are marked by *.

of Forestry and Technology (CSUFT). Axenic cultures were maintained in the China Forestry Culture Collection Centre (CFCC) and Central South University of Forestry and Technology Culture Collection (CSUFTCC).

Morphological and cultural characterization

Agar plugs (6 mm diam) were taken from the edge of actively growing cultures on PDA and transferred onto the centre of 9 cm diam Petri dishes containing 2% tap water agar supplemented with sterile pine needles (PNA) (Smith et al. 1996) and potato dextrose agar (PDA) and incubated at 25 °C under a 12 h near-ultraviolet light/12 h dark cycle to induce sporulation as described in recent studies (Gomes et al. 2013; Lombard et al. 2014). Colony characters and pigment production on PNA and PDA were noted in the 10-day culture. Colony features were rated according to the color charts of Rayner (1970). Cultures were examined periodically for the development of conidiomata. The microscopic examination was based on the morphological features of conidiomata obtained from the fungal growth mounted in clear lactic acid. At least 30 conidiomata and conidia were measured to calculate the mean size/length. Micro-morphological observations were done at ×1000 magnification using a Leica compound microscope (DM 2500) with interference contrast (DIC) optics. Descriptions, nomenclature, and illustrations of taxonomic novelties were deposited at MycoBank (www.MycoBank.org) (Crous et al. 2004).

DNA extraction, PCR amplification and sequencing

Genomic DNA was extracted from colonies grown on cellophane-covered PDA using a CTAB [cetyltrimethylammonium bromide] method (Doyle and Doyle 1990). For PCR amplifications of phylogenetic markers, five different primer pairs were used (Yang et al. 2018). The PCR conditions were: an initial denaturation step of 8 min at 95 °C followed by 35 cycles of 30 s at 95 °C, 30 s at 51 °C (ITS), 58 °C (*his3*) or 55 °C (*cal*, *tef-1a*, *tub2*), and 1 min at 72 °C, and a final elongation step of 5 min at 72 °C. PCR amplification products were assayed via electrophoresis in 2% agarose gels. DNA sequencing was performed using an ABI PRISM 3730XL DNA Analyzer with a BigDye Terminater Kit v.3.1 (Invitrogen, Waltham, MA, USA) at the Shanghai Invitrogen Biological Technology Company Limited (Beijing, China).

Phylogenetic analyses

The quality of our amplified nucleotide sequences was checked and combined by SeqMan v.7.1.0 and reference sequences (Table 1) were retrieved from the National Center for Biotechnology Information (NCBI), according to recent publications of the genus (Guo et al. 2020; Sun et al. 2021; Yang et al. 2021b). Sequences were aligned using MAFFT v. 6 (Katoh and Toh 2010) and manually corrected using Bioedit 7.0.9.0 (Hall 1999). Phylogenetic analyses were carried out with maximum likelihood analysis



Figure 1. Phylogram of *Diaporthe* resulting from a maximum likelihood analysis based on combined ITS, *cal*, *his3*, *tef-1a* and *tub2*. The tree is rooted with *Diaporthella corylina*. Values above the branches indicate Maximum Likelihood bootstrap (left, ML BP ≥ 75%) and Bayesian probabilities (right, BI PP ≥ 0.95). Strains in current study are in blue and the ex-type cultures are in bold.


Diaporthe betulae CFCC 50470
Diaporthe betulae CFCC 50469
Diaporthe padina CFCC 52591
Diaporthe padina CFCC 52590
Diaporthe betulina CFCC 52560
Diaporthe betulina CFCC 52561
Diaporthe chensiensis CFCC 52568
Diaporthe chensiensis CFCC 52567
Diaporthe vaccinii CBS 160.32
Diaporthe alleghaniensis CBS 495.72
Diaporthe eres (syn. D. celastrina) CBS 139.27
Diaporthe eres DAOM 695742
Diaporthe eres CGMCC 3.17081
Diaporthe eres MFLUCC 17-0963
Diaporthe eres AR5193
Diaporthe eres MFLU 17-0646
Diaporthe eres CFCC 52579
Diaporthe eres CFCC 52575
Diaporthe eres CGMCC 3.17089
Diaporthe eres (syn. D. camptothecicola) CFCC 51632
Diaporthe eres CFCC 52580
Diaporthe eres CFCC 52576
Diaporthe eres MFLUCC 16-0113
Diaporthe eres (syn. D. ellipticola) CGMCC 3.17084
Diaporthe eres (syn. D. celeris) CBS 143349
Diaporthe eres CFCC 52581
Diaporthe eres MAFF625034
Diaporthe eres CBS 587.79
Diaporthe eres CFCC 52578
Diaporthe eres CFCC 52577
Diaporthe eres CSUFTCC101
Diaporthe eres CSUFTCC102
Diaporthe eres CSUFTCC103
Diaporthe eres CGMCC 3.15181
Diaporthe eres CBS 338.89
Diaporthe eres AR5211
Diaporthe eres (syn. D. neilliae) CBS 144.27
Diaporthe eres (syn. D. alnea) CBS 146.46
Diaporthe virgiliae CMW40748
Diaporthe heterophyllae CPC 26215
Diaporthe grandiflori SAUCC194.84
Diaporthe penetriteum LC3215
Diaporthe collariana MFLU 17-2770
Diaporthe litchii SAUCC194.22
Diaporthe heliconiae SAUCC194.77
Diaporthe subclavata ZJUD95
Diaporthe tibetensis CFCC 52000
Diaporthe tibetensis CFCC 51999
Diaporthe coryli CFCC 53083
Diaporthe coryli CFCC 53084
Diaporthe ukurunduensis CFCC 52592
Diaporthe ukurunduensis CFCC 52593
Diaporthe citrichinensis CGMCC 3.15225
Diaporthe fraxinicola CFCC 52582
Diaporthe fraxinicola CFCC 52583
Diaporthe albosinensis CFCC 53067
Diaporthe albosinensis CFCC 53066
Diaporthe acerigena CFCC 52555
Diaporthe acerigena CFCC 52554
Diaporthe xunwuensis CFCC 53085
Diaporthe xunwuensis CFCC 53086
Diaporthe oraccinii LC3166
Diaporthe conica CFCC 52572
Diaporthe conica CFCC 52571
Diaporthe citri AR3405
Diaporthe citri CFCC 53079
Diaporthe apiculata CFCC 53068
Diaporthe apiculata CFCC 53069
Diaporthe charlesworthii BRIP 54884m
Diaporthe sennae CFCC 51637
Diaporthe sennae CFCC 51636
Diaporthe irregularis CGMCC 3.20092
Diaporthe bauhiniae CFCC 53072
Diaporthe bauhiniae CFCC 53071
Diaporthe aquatica IFRDCC 3051
Diaporthe psoraleae-pinnatae CPC 21638
Diaporthe shaanxiensis CFCC 53106
Diaporthe shaanxiensis CFCC 53107
Diaporthe incompleta LC6754
Diaporthe aspalathi CBS 117169
Diaporthe crotalariae CBS 162.33
Diaporthe caulivora CBS 127268
Diaporthe verniciicola CFCC 53110
Diaporthe verniciicola CFCC 53109
Diaporthe schisandrae CFCC 51989
Diaporthe schisandrae CFCC 51988
Diaporthe rostrata CFCC 50063
Diaporthe rostrata CFCC 50062
Diaporthe juglandicola CFCC 51134
Diaporthe minima CGMCC 3.20097
Diaporthe betulicola CFCC 51128
Diaporthe betulicola CFCC 51129
Diaporthe bohemiae CBS 143347
Diaporthe hispaniae CBS 143351
Diaporthe ampelina STE U2660
Diaporthe thunbergii MFLUCC 10-0576a

Figure 1. Continued.


Diaporthe amygdali (syn. D. garethjonesii) MFLUCC 12-0542a
Diaporthe amygdali (syn. D. chongqingensis) PSCG 435
Diaporthe amygdali (syn. D. ovoicicola) CGMCC 3.17093
Diaporthe amygdali (syn. D. terminalis) CGMCC 3.15183
Diaporthe amygdali (syn. D. fusicola) CGMCC 3.17087
Diaporthe amygdali (syn. D. kadsurae) CFCC 52587
Diaporthe amygdali (syn. D. kadsurae) CFCC 52586
Diaporthe amygdali CBS 126679
Diaporthe amygdali (syn. D. mediterranea) SAUCC194.111
Diaporthe amygdali (syn. D. sterilis) CBS 136969
Diaporthe celticola CFCC 53074
Diaporthe celticola CFCC 53075
Diaporthe celticola CFCC 53076
Diaporthe acaciigena CBS 129521
Diaporthe silvicola CFCC 54191
Diaporthe masirevii MFLU 17-2622
Diaporthe asheicola CBS 136967
Diaporthe australafricana CBS 111886
Diaporthe rudis AR3422
Diaporthe cynaroidis CBS 122676
Diaporthe taliciicola BRIP 54825
Diaporthe subcylindrospora KUMCC 170151
Diaporthe cissusis CAA 823
Diaporthe cassines CPC 21916
Diaporthe nothofagi BRIP 54801
Diaporthe benedicti SBen914
Diaporthe pseudorosicola MFLU 15-3228
Diaporthe acaciae CPC 26217
Diaporthe toulouensis MFLUCC 17-1051
Diaporthe tonkinensis CBS 534.93
Diaporthe spartinicola CPC 24951
Diaporthe pseudoalnea CFCC 54190
Diaporthe perjuncta CBS 109745
Diaporthe constrictospora CGMCC 3.20096
Diaporthe brasiliensis CBS 133183
Diaporthe caatingaensis URM7486
Diaporthe paranensis CBS 133184
Diaporthe malorum CAA 734
Diaporthe passiflorae CPC 19183
Diaporthe minusculata CGMCC 3.20098
Diaporthe eres CBS 133186
Diaporthe cinnamomi CFCC 52569
Diaporthe cinnamomi CFCC 52570
Diaporthe discoidispora ZJUD89
Diaporthe citriasiana CGMCC 3.15224
Diaporthe siamensis MFLUCC 10-0573a
Diaporthe yunnanensis LC6168
Diaporthe quercicola CSUFTCC104
Diaporthe quercicola CSUFTCC105
Diaporthe quercicola CSUFTCC106
Diaporthe eguttulata ZJUD47
Diaporthe passifloricola CPC 27480
Diaporthe durionigena VTCC 930005
Diaporthe thunbergicola MFLUCC 12-0033
Diaporthe rosae MFLUCC 17-2658
Diaporthe miriciae BRIP 54736j
Diaporthe ueckerae FAU656
Diaporthe unshiuensis CFCC 52594
Diaporthe unshiuensis CFCC 52595
Diaporthe longicolla FAU599
Diaporthe tectonendophytica MFLUCC 13-0471
Diaporthe masirevii BRIP 54256
Diaporthe kongii BRIP 54031
Diaporthe endophytica CBS 133811
Diaporthe fructicola MAFF 246408
Diaporthe subellipicola KUMCC 17-0153
Diaporthe heterostemmatis SAUCC194.85
Diaporthe kochmanii BRIP 54033
Diaporthe sojae FAU635
Diaporthe ovalispora ZJUD93
Diaporthe convolvuli CBS 124654
Diaporthe batatas CBS 122.21
Diaporthe melonis CBS 435.87
Diaporthe helianthi CBS 592.81
Diaporthe rosiphthora COAD 2914
Diaporthe schini CBS 133181
Diaporthe racemosae CPC 26646
Diaporthe terebinthifolii CBS 133180
Diaporthe gulyae BRIP 54025
Diaporthe angelicae CBS 111592
Diaporthe subordinaria CBS 464.90
Diaporthe cucurbitae DAOM 42078
Diaporthe guttulata CGMCC 3.20100
Diaporthe cichorii MFLUCC 17-1023
Diaporthe arctii DP0482
Diaporthe neoarctii CBS 109490
Diaporthe schoeni MFLU 15-1279
Diaporthe acericola MFLUCC 17-0956
Diaporthe novem CBS 127269
Diaporthe phaseolorum AR4203
Diaporthe lusitanicae CBS 123212
Diaporthe cuppatea CBS 117499
Diaporthe rossmaniae CAA 762
Diaporthe pyracanthae CAA 483
Diaporthe infecunda CBS 133812
Diaporthe serafiniae BRIP 55665a
Diaporthe beilharziae BRIP 54792
Diaporthe caryae CFCC 52564
Diaporthe caryae CFCC 52563
Diaporthe sackstonii BRIP 54669b
Diaporthe middletonii BRIP 54884e
Diaporthe myracrodruonis JRM 7972
Diaporthe arezzoensis MFLUCC 15-0127
Diaporthe sambucusii CFCC 51986
Diaporthe sambucusii CFCC 51987
Diaporthe compactum LC3083
Diaporthe ganjae CBS 180.91
Diaporthe alangii CFCC 52556
Diaporthe alangii CFCC 52557
Diaporthe hubeiensis JZB320123
Diaporthe tectonae MFLUCC 12-0777
Diaporthe tulliensis BRIP 62248a
Diaporthe sclerotioides CBS 296.67
Diaporthe longispora CBS 194.36
Diaporthe amygdalina CBS 114015
Diaporthe goulteri BRIP 55657a
Diaporthe raonikayaporum CBS 133182
Diaporthe neoraonikayaporum MFLUCC 14-1136
Diaporthe mayteni CBS 133185
Diaporthella corylina CBS 121124

Figure 1. Continued.

(ML), which was performed at the CIPRES web portal (Miller et al. 2010), 1000 rapid bootstrap replicates were run with GTRGAMMA model of nucleotide evolution. Bayesian inference analysis (BI) was performed in MrBayes v. 3.2.0 (Ronquist and Huelsenbeck 2003). The best-fit nucleotide substitution models for each gene were selected using jModelTest v. 2.1.7 (Darriba et al. 2012) under the Akaike Information Criterion. The best nucleotide substitution model for ITS, *his3* and *tub2* was TrN+I+G, while HKY+I+G was selected for both *cal* and *tef-1a*. Phylogenetic trees were viewed in FigTree v1.4. The names of the isolates from the present study are marked in blue in the trees. Maximum likelihood bootstrap support values ≥ 75% (BT) are given at the nodes. Bayesian posterior probabilities ≥ 0.95 (PP) were thickened in the phylogenetic tree. Alignment and trees were deposited in TreeBASE (submission ID: S29621).

Results

Phylogenetic analyses

The sequence datasets for the ITS, *cal*, *his3*, *tef-1a* and *tub2*, were analysed in combination to infer the interspecific relationships within *Diaporthe*. The combined species phylogeny of the *Diaporthe* isolates consisted of 303 sequences, including the outgroup *Diaporthella corylina* (CBS 121124). A total of 2535 characters including gaps (512 for ITS, 524 for *cal*, 525 for *his3*, 463 for *tef-1a*, and 511 for *tub2*) were included in the phylogenetic analysis. Similar tree topologies were obtained by ML and BI methods, and the best scoring ML tree is shown in Fig. 1. The ML analysis yielded a tree with a likelihood value of ln: -76822.498401 and the following model parameters: alpha: 0.508079; Π(A): 0.214617, Π(C): 0.326518, Π(G): 0.235187 and Π(T): 0.223678. The phylogenetic tree inferred from the concatenated alignment resolved the sixteen *Diaporthe* isolates from branch cankers or leaf spots into six well-supported monophyletic clades that represent four novel species and two known species of *Diaporthe* (Fig. 1).

Taxonomy

Diaporthe celticola **C.M. Tian & Q. Yang, sp. nov.**
MycoBank No: 832920
Fig. 2

Diagnosis. Distinguished from the other *Diaporthe* species based on DNA sequence data and characterised by conidiomata with single necks erumpent through the host bark.

Etymology. Named after the host genus on which it was collected, *Celtis*.

Description. ***Conidiomata*** pycnidial, 535–605 × 210–225 μm diam, solitary and with single necks erumpent through host bark. ***Ectostromatic disc*** brown, one ostiole per disc, with yellowish cream conidial drops exuding from the ostioles. Tissue around the neck is cylindrical. ***Locule*** circular, undivided, 350–375 μm diam. ***Conidiophores***

reduced to conidiogenous cells. ***Conidiogenous cells*** unbranched, straight or sinuous, apical or base sometimes swelling, (8–)10.5–13(–14.5) × 1–1.5 µm (n = 30), L/W = 8.5–10.5. ***Alpha conidia*** hyaline, aseptate, ellipsoidal, biguttulate, (5–)6–7 × 3.5–4 µm (n = 30), L/W = 1.5–1.8. ***Beta conidia*** not observed.

Culture characters. Colony originally flat with white fluffy aerial mycelium, becoming light brown to olive-green mycelium with age, marginal area irregularly, with yellowish cream conidial drops exuding from the ostioles.

Specimens examined. China, Zhejiang Province: Hanzhou City, on branches of *Celtis vandervoetiana*, 12 May 2018, *Q. Yang* & *Y.M. Liang* (holotype BJFC-S1616; ex-type living culture: CFCC 53074; living cultures: CFCC 53075 and CFCC 53076).

Notes. Three strains representing *Diaporthe celticola* cluster in a well-supported clade (ML/BI = 100/1), and appear closely related to *D. acaciigena*. *Diaporthe celticola* can be distinguished based on ITS, *cal*, *his3*, *tef-1a*, and *tub2* loci from *D. acaciigena* (29/473 in ITS, 68/442 in *cal*, 53/460 in *his3*, 79/330 in *tef-1a*, and 49/415 in *tub2*). Morphologically, *D. celticola* is characterised by conidiomata with single necks erumpent through the host bark and can be distinguished from *D. acaciigena* by smaller alpha conidia



Figure 2. *Diaporthe celticola* (BJFC-S1616) **a, b** habit of conidiomata on twig **c** transverse section through conidiomata **d, e** conidiogenous cells with alpha conidia **f** alpha conidia **g, h** conidiomata formed on PDA. Scale bars: 200 µm (**b, c**); 10 µm (**d–f**).

(6–7 × 3.5–4 vs. 10–11 × 6–6.5 μm) (Crous et al. 2011). This is the first occasion that *Diaporthe* species have been discovered from infected branches on *Celtis vandervoetiana* and demonstrates it to be a new species based on phylogeny and morphology.

Diaporthe eres **Nitschke, Pyrenomyc. Germ. 2: 245, 1870.**

Description. See Udayanga et al. (2014b).

Specimens examined. China. Beijing: Pinggu District, on branches of *Populus* × *xiaohei*, 10 July 2020, *Q. Yang* (CSUFT101; living cultures: CSUFTCC101, CSUFTCC102, and CSUFTCC103).

Notes. *Diaporthe eres* is the type species of the genus and was originally described by Nitschke (1870), from *Ulmus* sp. in Germany, which has a widespread distribution and a broad host range as an endophyte or saprobe, or pathogen causing leaf spots, stem cankers and diseases of woody plants (Udayanga et al. 2014b). In the present study, three isolates (CSUFTCC101, CSUFTCC102, and CSUFTCC103) are embedded into the *D. eres* species based on DNA sequence data (Fig. 1). We therefore describe *D. eres* as a known species for this clade.

Diaporthe meliae **C.M. Tian & Q. Yang, sp. nov.**

MycoBank No: 829523

Fig. 3

Diagnosis. Distinguished from the phylogenetically closely-related species, *D. podocarpi-macrophylli*, in shorter alpha conidia.

Etymology. Named after the host genus on which it was collected, *Melia*.

Description. ***Conidiomata*** pycnidial, immersed in the host bark, scattered, erumpent through the bark surface, discoid, with a single locule. ***Ectostromatic disc*** dark brown, one ostiole per disc, (325–)330–375(–385) μm (n = 30) diam. ***Locule*** undivided, 420–640 × 385–515 μm (n = 30). ***Conidiophores*** reduced to conidiogenous cells. ***Conidiogenous cells*** (13.5–)15–26.5(–28) × 1.3–2.1(–2.3) μm (n = 30), L/W = 8.5–15.5, cylindrical, hyaline, branched, straight or slightly curved, tapering towards the apex. ***Alpha conidia*** hyaline, aseptate, fusiform, multi-guttulate, (6.7–)8–9.5(–10) × (2–)2.1–2.3 μm (n = 30), L/W = 3.4–4.5. ***Beta conidia*** not observed.

Culture characters. Colony originally flat with white felty aerial mycelium, becoming auburn furcate mycelium with age, with irregular margin, conidiomata absent.

Specimens examined. China, Shandong Province: Rizhao City, on branches of *Melia azedarach*, 20 April 2018, *N. Jiang* (holotype BJFC-S1668; ex-type living culture: CFCC 53089; living culture: CFCC 53090).

Notes. Two strains representing *Diaporthe meliae* cluster in a well-supported clade (ML/BI = 100/1), and appear closely related to *D. podocarpi-macrophylli*. *Diaporthe meliae* can be distinguished based on ITS, *his3*, *tef-1a*, and *tub2* loci from *D. podocarpi-macrophylli* (4/459 in ITS, 15/455 in *his3*, 25/349 in *tef-1a*, and 14/401 in *tub2*).



Figure 3. *Diaporthe meliae* (BJFC-S1668) **a, b** habit of conidiomata on twig **c** transverse section through conidiomata **d** longitudinal section through conidiomata **e, f** conidiogenous cells **g** alpha conidia. Scale bars: 1 mm(**b**); 200 μm (**c, d**); 10 μm (**e–g**).

Morphologically, *D. meliae* can be distinguished from *D. podocarpi-macrophylli* by its longer conidiogenous cells (15–26.5 vs. 6–18 μm) and alpha conidia (8–9.5 vs. 3.5–8.5 μm) (Gao et al. 2017).

Diaporthe multiguttulata **F. Huang, K.D. Hyde & Hong Y. Li, Fungal Biology 119(5): 343, 2015.**
Fig. 4

Description. See Yang et al. (2021a).



Figure 4. *Diaporthe multiguttulata* (BJFC-S1615) **a, b** habit of conidiomata on twig **c** transverse section through conidiomata **d** longitudinal section through conidiomata **e–g** conidiomata formed on PDA **h** conidiogenous cells **i** alpha conidia **g** beta conidia. Scale bars: 200 μm (**b–d**); 10 μm (**h–j**).

Specimens examined. China, Jiangxi Province: Ganzhou City, on branches of *Citrus maxima*, 11 May 2018, *Q. Yang, & Y.M. Liang* (BJFC-S1615; living cultures: CFCC 53098, CFCC 53099, and CFCC 53100).

Notes. *Diaporthe multiguttulata* is characterised by ellipsoidal alpha conidia with one large guttulate, and was originally described as an endophyte from healthy branch of *Citrus grandis* in Fujian Province, China (Huang et al. 2015). Yang et al. (2021a) identified three isolates from *Citrus maxima* as *D. multiguttulata* based on DNA sequence data and confirmed from the morphological characters. In the present study, isolates (CFCC 53098, CFCC 53099, and CFCC 53100) from an additional specimen were observed and supplemented with beta conidia (Fig. 4j).

Diaporthe quercicola **Q. Yang, sp. nov.**

MycoBank No: 843494

Fig. 5

Diagnosis. Distinguished from the phylogenetically closely-related species, *D. biguttulata*, by its filiform, eguttulate alpha conidia.

Etymology. Named after the host genus on which it was collected, *Quercus*.



Figure 5. *Diaporthe quercicola* (CSUFTCC104) **a** conidiomata formed on PDA **b** conidiogenous cells **c** alpha and beta conidia. Scale bars: 200 μm (**a**); 10 μm (**b, c**).

Description. On PDA: ***Conidiomata*** pycnidial, 250–330 μm diam, globose, solitary or aggregated, deeply embedded in the medium, erumpent, single or clustered in groups of 3–5 pycnidia, coated with hyphae, cream to yellowish translucent conidial droplets exuded from the ostioles. ***Conidiophores*** reduced to conidiogenous cells. ***Conidiogenous cells*** hyaline, cylindrical, unbranched, straight, tapering towards the apex, (17–)20–26(–34.5) × 2.5–3.5 μm (n = 30), L/W = 6.5–9. ***Alpha conidia*** (6.5–)7–8.5(–9) × (1.5–)2–3 μm (n = 30), L/W = 3–4.5, aseptate, hyaline, fusiform, apex at both ends, eguttulate. ***Beta conidia*** hyaline, aseptate, filiform, straight or sinuous at one end, eguttulate, (21.5–)25.5–31(–33) × 1 μm (n = 30), L/W = 22.5–31.5.

Culture characters. Colony at first white, becoming dark brown with age. Aerial mycelium white, dense, fluffy, with yellowish conidial drops exuding from the ostioles after 20 days.

Specimens examined. China. Shaanxi Province: Xian City, on branches of *Quercus aliena*, 10 July 2020, *Q. Yang* (holotype CSUFTCC104; ex-type living culture: CSUFTCC104; living cultures: CSUFTCC105 and CSUFTCC106).

Notes. Three strains representing *Diaporthe quercicola* cluster in a well-supported clade (ML/BI = 100/1), and appear closely related to *D. biguttulata*. *Diaporthe quercicola* can be distinguished based on ITS, *his3*, and *tef-1a* loci from *D. biguttulata* (8/461 in ITS, 18/448 in *his3*, and 22/325 in *tef-1a*). Morphologically, *D. quercicola* can be distinguished from *D. biguttulata* by its fusiform, eguttulate alpha conidia and narrower beta conidia (1 vs. 0.9–1.6 μm) (Huang et al. 2015).

Diaporthe rhodomyrti **C.M. Tian & Q. Yang, sp. nov.**
MycoBank No: 829525
Fig. 6

Diagnosis. Distinguished from the phylogenetically closely-related species, *D. hongkongensis*, in narrower beta conidia.



Figure 6. *Diaporthe rhodomyrti* (BJFC-S1660) **a** conidioma formed on PNA **b** conidiogenous cells **c** alpha and beta conidia. Scale bars: 500 μm (**a**); 10 μm (**b, c**).

Etymology. Named after the host genus on which it was collected, *Rhodomyrtus*.

Description. On PNA: ***Conidiomata*** pycnidial, 500–850 μm diam, globose or rostrate, black, erumpent in tissue, erumpent at maturity, often with translucent conidial drops exuding from ostioles. ***Conidiophores*** reduced to conidiogenous cells. ***Conidiogenous cells*** (14.5–)15.5–23(–25.5) × 1.5–2 μm (n = 30), L/W = 8.5–13, cylindrical, hyaline, unbranched, septate, straight, tapering towards the apex. ***Alpha conidia*** abundant in culture, hyaline, aseptate, ellipsoidal, biguttulate, 6–7(–8.5) × 2–2.5(–3) μm (n = 30), L/W = 2.8–3.3. ***Beta conidia*** hyaline, aseptate, filiform, straight to sinuous, eguttulate, (15–)16.5–21.5(–23) × 1–1.5 μm (n = 30), L/W = 15.5–16.5.

Culture characters. Colony entirely white at surface, reverse with pale brown pigmentation, white, fluffy aerial mycelium.

Specimens examined. China. Jiangxi Province: Ganzhou City, on leaves of *Rhodomyrtus tomentosa*, 10 May 2018, *Q. Yang* & *Y.M. Liang* (holotype BJFC-S1660; ex-type living culture: CFCC 53101; living culture: CFCC 53102).

Notes. This new species is introduced as molecular data, and shows it to be a distinct clade with high support (ML/BI = 100/1) and appears closely related to *Diaporthe hongkongensis*. *Diaporthe rhodomyrti* can be distinguished based on ITS, *cal*, *his3*, *tef-1a*, and *tub2* loci from *D. hongkongensis* (2/463 in ITS, 26/441 in *cal*, 11/434 in *his3*, 10/327 in *tef-1a*, and 2/420 in *tub2*). Morphologically, *D. rhodomyrti* can be distinguished from *D. hongkongensis* by its longer conidiogenous cells (15.5–23 vs. 5–12 μm) and narrower beta conidia (1–1.5 vs. 1.5–2 μm) (Gomes et al. 2013). This is the first time that *Diaporthe* species has been discovered from infected leaves on *Rhodomyrtus tomentosa* and demonstrate it as a new species based on phylogeny and morphology.

Discussion

In this study, investigations of forest pathogens in Beijing, Jiangxi, Shaanxi and Zhejiang Provinces was carried out. Identification of our collections was conducted, based on isolates from fruiting bodies using five combined loci (ITS, *cal*, *his3*, *tef-1a*, and *tub2*), as well as morphological characteristics. It includes *Diaporthe eres* and

D. multiguttulata, as well as four new species named *D. celticola*, *D. meliae*, *D. quercicola*, and *D. rhodomyrti*.

Diaporthe (Diaporthaceae, Sordariomycetes) are species-rich asexual taxa, which are common pathogens that cause a variety of diseases, including dieback, stem cankers, leaf spots, leaf and pod blights, fruit rots and seed decay (Uecker 1988; Rehner and Uecker 1994; Mostert et al. 2001; Thompson et al. 2001; Santos et al. 2011). Because many *Diaporthe* species have overlapping morphological traits, sequence data is essential to resolve this genus and introduce new species (Udayanga et al. 2014a). Combined gene sequence of ITS, *cal*, *his3*, *tef-1a*, and *tub2* is the optimal combination for species delimitation (Santos et al. 2017). However, removing the ITS locus has little effect on reconstructed phylogenies, identifying the *cal-his3-tef-1a-tub2* four loci tree as almost equivalent to the five loci phylogenetic tree.

Many confusions occur in species separation of *Diaporthe eres* complex with the lack of an ex-type culture or ex-epitype culture, although a broad species concept has historically been associated with *D. eres* (Udayanga et al. 2014b). Fan et al. (2018) demonstrated the effectiveness of three loci, including *cal*, *tef-1a* and *tub2*, for the identification of the *D. eres* complex in walnut trees. Similarly, Yang et al. (2018) also used three-locus sequences to identify *D. eres* species associated with different hosts in China, and Chaisiri et al. (2021) revealed the phylogenetic analysis from the combined dataset of *cal*, *his3*, *tef-1a* and *tub2* was highly effective, but the ITS region impeded species delimitation, which conforms with Yang et al. (2018).

Recently, several studies have been conducted associated with various hosts in China. For instance, the research conducted by Guo et al. (2020) revealed six novel *Diaporthe* species that infect pears and are responsible for pear shoot canker. Sun et al. (2021) showed high species diversity of *Diaporthe* in tropical rain forests, with description of eight new species. Wang et al. (2021) represented the first characterization of *Diaporthe* species associated with peach constriction canker in China, and contributed useful data for practicable disease management. Yang et al. (2021b) identified two new species from *Camellia oleifera*, which is an important edible oil woody plant in southern China. This study also characterises the taxonomic and morphological diversity of *Diaporthe* spp. associated with different hosts, which indicated there is a potential of *Diaporthe* species remains to be discovered in China.

Acknowledgements

This study is financed by National Natural Science Foundation of China (Project No.: 31670647) and the Research Foundation of Education Bureau of Hunan Province (Project No.: 19B608).

References

Barr ME (1978) The Diaporthales in North America with emphasis on *Gnomonia* and its segregates. Mycologia Memoir 7: 1–232.

Castlebury LA, Rossman AY, Jaklitsch WJ, Vasilyeva LN (2002) A preliminary overview of the Diaporthales based on large subunit nuclear ribosomal DNA sequences. Mycologia 94(6): 1017–1031. https://doi.org/10.1080/15572536.2003.11833157

Chaisiri C, Liu X, Lin Y, Fu Y, Zhu F, Luo C (2021) Phylogenetic and haplotype network analyses of *Diaporthe eres* species in China based on sequences of multiple loci. Biology 10(3): e179. https://doi.org/10.3390/biology10030179

Crous PW, Gams W, Stalpers JA, Robert V, Stegehuis G (2004) MycoBank: An online initiative to launch mycology into the 21st century. Studies in Mycology 50: 19–22.

Crous PW, Groenewald JZ, Shivas RG, Edwards J, Seifert KA, Alfenas AC, Alfenas RF, Burgess TI, Carnegie AJ, Hardy GESJ, Hiscock N, Hüberli D, Jung T, Louis-Seize G, Okada G, Pereira OL, Stukely MJC, Wang W, White GP, Young AJ, McTaggart AR, Pascoe IG, Porter IJ, Quaedvlieg W (2011) Fungal Planet description sheets: 69–91. Persoonia 26(1): 108–156. https://doi.org/10.3767/003158511X581723

Darriba D, Taboada GL, Doallo R, Posada D (2012) jModelTest 2: More models, new heuristics and parallel computing. Nature Methods 9(8): e772. https://doi.org/10.1038/nmeth.2109

Dissanayake AJ, Zhang W, Liu M, Hyde KD, Zhao WS, Li XH, Yan JY (2017) *Diaporthe* species associated with peach tree dieback in Hubei, China. Mycosphere 8(5): 533–549. https://doi.org/10.5943/mycosphere/8/5/2

Doyle JJ, Doyle JL (1990) Isolation of plant DNA from fresh tissue. Focus 12: 13–15. https://doi.org/10.2307/2419362

Du Z, Fan XL, Hyde KD, Yang Q, Liang YM, Tian CM (2016) Phylogeny and morphology reveal two new species of *Diaporthe* from *Betula* spp. in China. Phytotaxa 269(2): 90–102. https://doi.org/10.11646/phytotaxa.269.2.2

Fan XL, Hyde KD, Udayanga D, Wu XY, Tian CM (2015) *Diaporthe rostrata*, a novel ascomycete from *Juglans mandshurica* associated with walnut dieback. Mycological Progress 14(10): 1–8. https://doi.org/10.1007/s11557-015-1104-5

Fan XL, Yang Q, Bezerra JDP, Alvarez LV, Tian CM (2018) *Diaporthe* from walnut tree (*Juglans regia*) in China, with insight of *Diaporthe eres* complex. Mycological Progress 17(7): 1–13. https://doi.org/10.1007/s11557-018-1395-4

Gao YH, Liu F, Duan W, Crous PW, Cai L (2017) *Diaporthe* is paraphyletic. IMA Fungus 8(1): 153–187. https://doi.org/10.5598/imafungus.2017.08.01.11

Gomes RR, Glienke C, Videira SIR, Lombard L, Groenewald JZ, Crous PW (2013) *Diaporthe*: A genus of endophytic, saprobic and plant pathogenic fungi. Persoonia 31(1): 1–41. https://doi.org/10.3767/003158513X666844

Guarnaccia V, Crous PW (2017) Emerging citrus diseases in Europe caused by species of *Diaporthe*. IMA Fungus 8(2): 317–334. https://doi.org/10.5598/imafungus.2017.08.02.07

Guarnaccia V, Crous PW (2018) Species of *Diaporthe* on *Camellia* and *Citrus* in the Azores Islands. Phytopathologia Mediterranea 57: 307–319. https://doi.org/10.14601/Phytopathol_Mediterr-23254

Guarnaccia V, Groenewald JZ, Woodhall J, Armengol J, Cinelli T, Eichmeier A, Ezra D, Fontaine F, Gramaje D, Gutierrez-Aguirregabiria A, Kaliterna J, Kiss L, Larignon P, Luque J,

Mugnai L, Naor V, Raposo R, Sándor E, Váczy KZ, Crous PW (2018) *Diaporthe* diversity and pathogenicity revealed from a broad survey of grapevine diseases in Europe. Persoonia 40(1): 135–153. https://doi.org/10.3767/persoonia.2018.40.06

Guo YS, Crous PW, Bai Q, Fu M, Yang MM, Wang XH, Du YM, Hong N, Xu WX, Wang GP (2020) High diversity of *Diaporthe* species associated with pear shoot canker in China. Persoonia 45(1): 132–162. https://doi.org/10.3767/persoonia.2020.45.05

Hall T (1999) BioEdit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. Nucleic Acids Symposium Series 41: 95–98.

Huang F, Udayanga D, Wang X, Hou X, Mei X, Fu Y, Hyde KD, Li HY (2015) Endophytic *Diaporthe* associated with *Citrus*: A phylogenetic reassessment with seven new species from China. Fungal Biology 119(5): 331–347. https://doi.org/10.1016/j.funbio.2015.02.006

Huang ST, Xia JW, Zhang XG, Sun WX (2021) Morphological and phylogenetic analyses reveal three new species of *Diaporthe* from Yunnan, China. MycoKeys 78: 49–77. https://doi.org/10.3897/mycokeys.78.60878

Jiang N, Voglmayr H, Piao CG, Li Y (2021) Two new species of *Diaporthe* (Diaporthaceae, Diaporthales) associated with tree cankers in the Netherlands. MycoKeys 85: 31–56. https://doi.org/10.3897/mycokeys.85.73107

Katoh K, Toh H (2010) Parallelization of the MAFFT multiple sequence alignment program. Bioinformatics 26(15): 1899–1900. https://doi.org/10.1093/bioinformatics/btq224

Lombard L, Van Leeuwen GCM, Guarnaccia V, Polizzi G, Van Rijswick PC, Karin Rosendahl KC, Gabler J, Crous PW (2014) *Diaporthe* species associated with *Vaccinium*, with specific reference to Europe. Phytopathologia Mediterranea 53: 287–299. https://doi.org/10.14601/Phytopathol_Mediterr-14034

Manawasinghe IS, Dissanayake AJ, Li X, Liu M, Wanasinghe DN, Xu J, Zhao W, Zhang W, Zhou Y, Hyde KD, Brooks S, Yan J (2019) High genetic diversity and species complexity of *Diaporthe* associated with grapevine dieback in China. Frontiers in Microbiology 10: e1936. https://doi.org/10.3389/fmicb.2019.01936

Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for Inference of Large Phylogenetic Trees. Institute of Electrical and Electronics Engineers: New Orleans, LA, USA. https://doi.org/10.1109/GCE.2010.5676129

Mostert L, Crous PW, Kang JC, Phillips AJ (2001) Species of *Phomopsis* and a *Libertella* sp. occurring on grapevines with specific reference to South Africa: Morphological, cultural, molecular and pathological characterization. Mycologia 93(1): 146–167. https://doi.org/10.1080/00275514.2001.12061286

Muralli TS, Suryanarayanan TS, Geeta R (2006) Endophytic *Phomopsis* species: Host range and implications for diversity estimates. Canadian Journal of Microbiology 52(7): 673–680. https://doi.org/10.1139/w06-020

Nitschke T (1870) Pyrenomycetes Germanici. Breslau. Eduard Trewendt, Germany. 2: e245.

Rayner RW (1970) A mycological colour chart. Commonwealth Mycological Institute, Kew, UK.

Rehner SA, Uecker FA (1994) Nuclear ribosomal internal transcribed spacer phylogeny and host diversity in the coelomycete *Phomopsis*. Canadian Journal of Botany 72(11): 1666–1674. https://doi.org/10.1139/b94-204

Ronquist F, Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. Bioinformatics 19(12): 1572–1574. https://doi.org/10.1093/bioinformatics/btg180

Rossman AY, Farr DF, Castlebury LA (2007) A review of the phylogeny and biology of the Diaporthales. Mycoscience 48(3): 135–144. https://doi.org/10.1007/S10267-007-0347-7

Santos JM, Phillips AJL (2009) Resolving the complex of *Diaporthe* (*Phomopsis*) species occurring on *Foeniculum vulgare* in Portugal. Fungal Diversity 34: 111–125.

Santos JM, Vrandečić K, Ćosić J, Duvnjak T, Phillips AJL (2011) Resolving the *Diaporthe* species occurring on soybean in Croatia. Persoonia 27(1): 9–19. https://doi.org/10.3767/003158511X603719

Santos L, Alves A, Alves R (2017) Evaluating multi-locus phylogenies for species boundaries determination in the genus *Diaporthe*. PeerJ 5: e3120. https://doi.org/10.7717/peerj.3120

Smith H, Wingfeld MJ, Coutinho TA, Crous PW (1996) *Sphaeropsis sapinea* and *Botryosphaeria dothidea* endophytic in *Pinus* spp. and *Eucalyptus* spp. in South Africa. South African Journal of Botany 62(2): 86–88. https://doi.org/10.1016/S0254-6299(15)30596-2

Sun W, Huang S, Xia J, Zhang X, Li Z (2021) Morphological and molecular identification of *Diaporthe* species in south-western China, with description of eight new species. MycoKeys 77: 65–95. https://doi.org/10.3897/mycokeys.77.59852

Thompson SM, Tan YP, Young AJ, Neate SM, Aitken EAB, Shivas RG (2001) Stem cankers on sunflower (*Helianthus annuus*) in Australia reveal a complex of pathogenic *Diaporthe* (*Phomopsis*) species. Persoonia 27(1): 80–89. https://doi.org/10.3767/003158511X617110

Udayanga D, Liu X, McKenzie EH, Chukeatirote E, Bahkali AH, Hyde KD (2011) The genus *Phomopsis*: Biology, applications, species concepts and names of common phytopathogens. Fungal Diversity 50(1): 189–225. https://doi.org/10.1007/s13225-011-0126-9

Udayanga D, Castlebury LA, Rossman AY, Hyde KD (2014a) Species limits in *Diaporthe*: Molecular re-assessment of *D. citri*, *D. cytosporella*, *D. foeniculina* and *D. rudis*. Persoonia 32(1): 83–101. https://doi.org/10.3767/003158514X679984

Udayanga D, Castlebury LA, Rossman AY, Chukeatirote E, Hyde KD (2014b) Insights into the genus *Diaporthe*: Phylogenetic species delimitation in the *D. eres* species complex. Fungal Diversity 67(1): 203–229. https://doi.org/10.1007/s13225-014-0297-2

Udayanga D, Castlebury LA, Rossman AY, Chukeatirote E, Hyde KD (2015) The *Diaporthe sojae* species complex: Phylogenetic re-assessment of pathogens associated with soybean, cucurbits and other field crops. Fungal Biology 119(5): 383–407. https://doi.org/10.1016/j.funbio.2014.10.009

Uecker FA (1988) A world list of *Phomopsis* names with notes on nomenclature, morphology and biology. Mycological Memoirs 13: 1–231.

von Höhnel FXR (1917) System der Diaportheen. Borntraeger.

Wang X, Guo Y, Du Y, Yang Z, Huang X, Hong N, Xu WX, Wang GP (2021) Characterization of *Diaporthe* species associated with peach constriction canker, with two novel species from China. MycoKeys 80: 77–90. https://doi.org/10.3897/mycokeys.80.63816

Wehmeyer LE (1975) The pyrenomycetous fungi. Mycologia Memoir 6: 1–250.

Yang Q, Fan XL, Du Z, Tian CM (2017) *Diaporthe* species occurring on *Senna bicapsularis* in southern China, with descriptions of two new species. Phytotaxa 302(2): 145–155. https://doi.org/10.11646/phytotaxa.302.2.4

Yang Q, Fan XL, Guarnaccia V, Tian CM (2018) High diversity of *Diaporthe* species associated with dieback diseases in China, with twelve new species described. MycoKeys 39: 97–149. https://doi.org/10.3897/mycokeys.39.26914

Yang Q, Jiang N, Tian CM (2020) Three new *Diaporthe* species from Shaanxi Province, China. MycoKeys 67: 1–18. https://doi.org/10.3897/mycokeys.67.49483

Yang Q, Jiang N, Tian CM (2021a) New species and records of *Diaporthe* from Jiangxi Province, China. MycoKeys 77: 41–64. https://doi.org/10.3897/mycokeys.77.59999

Yang Q, Tang J, Zhou GY (2021b) Characterization of *Diaporthe* species on *Camellia oleifera* in Hunan Province, with descriptions of two new species. MycoKeys 84: 15–33. https://doi.org/10.3897/mycokeys.84.71701